1837-1838 Caspar Morris [redacted] medical notes 1999

and is thrown off but a long continued use in small
Doses may be introduced into the System and
produce Colic Pictons & general pain in combination
with opium. Dr Wood mentions once had the use of
Nitrate of Silver suspended the pain to prevent its being
decomposed. Frequently acid externally & sedatives
internally may be reversed. Dr W has met with a
case. Opium to be given in cases of over Dose
Sulph Soda. Most acid in Hemorrhages from lungs
and uterus. They some sedative influence no
stimulating properties. May be given every
hour Euclin drops & Dialy Tinc Dose 1/10 gr
Every hour or Two. but not to be too long taken
lest it poisons. also applicable to all hemorrhages
Dr W used it in case of Bleeding from Bladder or
Kidneys: prompt in its action here. Marked Benefit
in Diarrhoea whether acute or Chronic. Combined
with Opium Ipecac & Calomel. White Stools
like Rice water. here use Cal & acetate
Dysentery maybe added Emetics useful
in bleeding of Phthisis. useful in Colliquative Diarrhoea
sometimes Dangerous antiphlogistic in its Effects on the
Inflammation of Mucous membrane of the Stomach. In
the latter stages of Yellow Fever there is a Subacute
inflammation of Stomach with Vomiting leading to
Black Vomit. here it may used with advantage
Dr Irving of Charleston. Dr W recommends its combination
these cases Dose 10 Drops every hour or Two. Kalchen
used also in Nervous Diseases. May be used
in Nervous cases where there is Mr Agitation